

PATENT COOPERATION TREATY

From the INTERNATIONAL BUREAU

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

To:

Assistant Commissioner for Patents
 United States Patent and Trademark
 Office
 Box PCT
 Washington, D.C. 20231
 ETATS-UNIS D'AMERIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 29 May 2000 (29.05.00)	
International application No. PCT/US99/25350	Applicant's or agent's file reference WSUR-1-14524
International filing date (day/month/year) 26 October 1999 (26.10.99)	Priority date (day/month/year) 27 October 1998 (27.10.98)
Applicant KAMMERECK, Rudolf et al	

1. The designated Office is hereby notified of its election made:

☒

in the demand filed with the International Preliminary Examining Authority on:

28 April 2000 (28.04.00)

☐

in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was☐

was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32:2(b).

The International Bureau of WIPO
 34, chemin des Colombettes
 1211 Geneva 20, Switzerland

Facsimile No. (41 22) 740 14 35

Authorized officer

Pascal Piriou

Telephone No. (41 22) 228 92 22

PATENT COOPERATION TREATY

PCT

REC'D 29 JAN 2001

INTERNATIONAL PRELIMINARY EXAMINATION REPORT PCT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference WSUR-1-14524	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/US99/25350	International filing date (day/month/year) 26 OCTOBER 1999	Priority date (day/month/year) 27 OCTOBER 1998
International Patent Classification (IPC) or national classification and IPC Please See Supplemental Sheet.		
Applicant WASHINGTON STATE UNIVERSITY RESEARCH FOUNDATION		

- This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.
- This REPORT consists of a total of 4 sheets.
☐ This report is also accompanied by ANNEXES, i.e., sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority. (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).
 These annexes consist of a total of 0 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☐ Non-establishment of report with regard to novelty, inventive step or industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☐ Certain observations on the international application

Date of submission of the demand 28 APRIL 2000	Date of completion of this report 18 DECEMBER 2000
Name and mailing address of the IPEA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer HELEN PRATT <i>[Signature]</i>
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0651

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/25350

I. Basis of the report**1. With regard to the elements of the international application: ***

- ☐ the international application as originally filed
- ☒ the description:
pages 1-27 , as originally filed
pages NONE , filed with the demand
pages NONE , filed with the letter of _____
- ☒ the claims:
pages 28-32 , as originally filed
pages NONE , as amended (together with any statement) under Article 19
pages NONE , filed with the demand
pages NONE , filed with the letter of _____
- ☒ the drawings:
pages NONE , as originally filed
pages NONE , filed with the demand
pages NONE , filed with the letter of _____
- ☒ the sequence listing part of the description:
pages NONE , as originally filed
pages NONE , filed with the demand
pages NONE , filed with the letter of _____

2. With regard to the language, all the elements marked above were available or furnished to this Authority in the language in which the international application was filed, unless otherwise indicated under this item.

These elements were available or furnished to this Authority in the following language _____ which is:

- ☐ the language of a translation furnished for the purposes of international search (under Rule 23.1(b)).
- ☐ the language of publication of the international application (under Rule 48.3(b)).
- ☐ the language of the translation furnished for the purposes of international preliminary examination (under Rules 55.2 and/or 55.3).

3. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international preliminary examination was carried out on the basis of the sequence listing:

- ☐ contained in the international application in printed form.
- ☐ filed together with the international application in computer readable form.
- ☐ furnished subsequently to this Authority in written form.
- ☐ furnished subsequently to this Authority in computer readable form.
- ☐ The statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.
- ☐ The statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished.

4. ☒ The amendments have resulted in the cancellation of:

- ☒ the description, pages NONE
- ☒ the claims, Nos. NONE
- ☒ the drawings, sheets/fig. NONE

5. ☐ This report has been drawn as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed, as indicated in the Supplemental Box (Rule 70.2(c)).**

* Replacement sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to this report since they do not contain amendments (Rules 70.16 and 70.17).

**Any replacement sheet containing such amendments must be referred to under item 1 and annexed to this report.

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No.

PCT/US99/25350

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

1. statement

Novelty (N)	Claims <u>1-41</u>	YES
	Claims <u>none</u>	NO
Inventive Step (IS)	Claims <u>none</u>	YES
	Claims <u>1-41</u>	NO
Industrial Applicability (IA)	Claims <u>1-41</u>	YES
	Claims <u>none</u>	NO

2. citations and explanations (Rule 70.7)

Claims 1-41 lack an inventive step under PCT Article 33(3) as being obvious over Dutton in view of Gunnerson et al. and Yamagishi.

Dutton discloses a plant protective coating composition containing ammonium linoleate in water, bentonite, and paraffin wax and aluminum flakes (page 2, lines 10-45). Claims 1 and 2 differ from the reference in that the composition is used to coat vegetables and fruits. Gunnerson et al. disclose that it is known to coat fruit with wax in vegetable oil and protein (abstract). Yamagishi disclose a composition containing paraffin, stearic acid and water and morpholine which can be sprayed on seedlings and saplings. As the various ingredients are known for coating of vegetable material, it would have been obvious to choose the food safe materials to coat fruits and vegetables for the function of protecting them from fungus and tree scald.

Claims 3, 8, 9 further require an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier. Dutton discloses the use of ammonium salts of unsaturated fatty acids (page 2, lines 36-54). Yamagishi discloses the use of morpholine, triethanolamine and higher fatty acids (abstract). Therefore, it would have been obvious to use known emulsifiers for their known functions.

Claims 4, 6 and 7 require particular waxes. Dutton discloses the use of paraffin, carnauba and beeswax (page 2, col. 1, lines 36-50). Therefore, it would have been obvious to use known ingredients for their known functions in making the coating.

Claims 5 and 10 require various proportions of clay to wax emulsion and claim 10 various amounts of dilution. The reference to Dutton shows a much higher dilution with water (page 2, col. 2, lines 10-25). However, the claims are not limited to the particular ingredients and particularly as in claim 5 water could be added in various amounts. Therefore, it would have been obvious to dilute the composition of clay and wax emulsion, to (Continued on Supplemental Sheet.)

Supplemental Box

(To be used when the space in any of the preceding boxes is not sufficient)

Continuation of: Boxes I - VIII

Sheet 10

CLASSIFICATION:

The International Patent Classification (IPC) and/or the National classification are as listed below:

IPC(7): A23B 4/10, A23L 1/059; A01G 13/00 and US Cl.: 47/ Digest 6 and Digest 11; 426/99, 102, 615

V. 2. REASONED STATEMENTS - CITATIONS AND EXPLANATIONS (Continued):

whatever dilution which would have the claimed effect.

The limitations of claims 11-23 have been discussed above and are obvious for those reasons, particularly as Dutton is to a plant protective composition (page 1, col. 1, lines 1-5). Therefore, it would have been obvious to use the composition of the combined references to protect plants.

Claim 24 further requires spraying the composition (claim 24) onto a plant, in particular amounts as in claim 25, multiple times as in claim 26. Dutton discloses diluting the composition for use in spraying. It would have been within the skill of the ordinary worker to dilute a known composition used for its known function to particular effective amounts. Therefore, it would have been obvious to spray the product in particular amounts multiple times.

Claim 27 further requires that the claimed composition is used to control insects. Dutton discloses that his composition can be used against borer attacks (page 3, col. 1, lines 5-10). The limitations of claims 28-41 have been discussed above. Therefore, it would have been obvious to use the claimed composition against insect attacks.

Claims 1-41 meets the requirements for industrial applicability as defined by PCT Article 33(4) in providing a plant protective coating and a method of protecting plants from insect damage and sunburn.

----- NEW CITATIONS -----

NONE



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A23B 4/10, A23L 1/059, A01G 13/00	A1	(11) International Publication Number: WO 00/24264 (43) International Publication Date: 4 May 2000 (04.05.00)
(21) International Application Number: PCT/US99/25350 (22) International Filing Date: 26 October 1999 (26.10.99) (30) Priority Data: 60/106,059 27 October 1998 (27.10.98) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US Not furnished (CIP) Filed on Not furnished (71) Applicant (for all designated States except US): WASHINGTON STATE UNIVERSITY RESEARCH FOUNDATION [US/US]; N.E. 1610 Eastgate Boulevard, Pullman, WA 99164-1802 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): KAMMERECK, Rudolf [US/US]; 1129 Eighth Street, Wenatchee, WA 98801 (US). SCHRADER, Lawrence, E. [US/US]; 3504 Crestview Street, Wenatchee, WA 98801 (US).		(74) Agent: BRADLEY, R., Douglas; Christensen O'Connor Johnson & Kindness PLLC, Suite 2800, 1420 Fifth Avenue, Seattle, WA 98101 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: USE OF CLAY AND LIPID FORMULATIONS TO PROTECT HORTICULTURAL CROPS FROM SUNBURN AND INSECT DAMAGE		
(57) Abstract <p>Sunburn and insect damage to fruit and vegetable crops is significantly reduced by treatment of both fruit and foliage with a preventative amount of thixotropic smectic clay material, chemically altered to render its surface lipophilic, which is combined with a wax emulsion comprising a matrix of complex hydrocarbons, an emulsifying agent and water. In the practice of this invention the sunburn and insect protective composition is further diluted in an aqueous solution that is sprayable by commercial applicators.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

USE OF CLAY AND LIPID FORMULATIONS TO PROTECT HORTICULTURAL CROPS FROM SUNBURN AND INSECT DAMAGE

Field of the Invention

The invention relates to protective coated fruits and vegetables, and methods
5 for the treatment of plants that reduces the incidence of insect and sunburn damage.

Background of the Invention

Sunburn has been a problem for apple growers for at least 75 years, but its
incidence has increased in recent years with the widespread use of dwarfing
rootstocks and high-density plantings. Many cultivars (e.g., 'Fuji,' 'Granny Smith,'
10 'Jonagold,' 'Gala,' and 'Braeburn') are susceptible to sunburn. Prominent growers
have indicated that sunburn may be the most significant cullage or quality problem in
the industry. Trees are smaller and fruit are more exposed to solar radiation making
fruit more susceptible to sunburn.

There is no adequate product on the market today for preventing sunburn
15 damage. Many growers use overhead evaporative cooling or shade cloth to reduce
sunburn in their apple orchards. Evaporative cooling decreases the temperature of
the fruit and helps protect the fruit from sunburn (Parchomchuk, P. and Meheriuk,
M., "Orchard cooling With Pulsed Overtree Irrigation to Prevent Solar Injury and
Improve Fruit Quality of 'Jonagold' Apples," *HortScience* 31:802-804 (1996)).
20 However, growers are concerned about several deleterious effects on fruit trees and
soil (Warner, G., "Overhead Cooling May Not Be Total Sunburn Cure," *Good Fruit
Grower* 46(12):20-21 (1995)). The shade cloths cost several thousand dollars per acre
to install, and frequently interfere with normal color development of fruit. Uniform

shade causes an undesirable alteration in the growth habit of apple trees and significantly reduces fruit production (Warner, G., "Cooling Problems Prompt Growers To Try Covers," *Good Fruit Grower* 46(12):24-25 (1995); Warner, G., "Growers Test Shade Cloths To Reduce Fuji Sunburn," *Good Fruit Grower* 46(17):55-63 (1995); Warner, G., "What Shade Do Cloths Provide, What Do You Need?", *Good Fruit Grower* 46(17):50-53 (1995)). Problems with these approaches confirm that new treatments are needed to lower fruit temperature, but not interfere with color development or fruit production.

In 1986 and 1987, Sibbett et al. ("Effect Of A Topically Applied Whitener On Sun Damage To Granny Smith Apples," *California Agriculture* 45(1):9-10 (1991)) in California attempted to solve the problem by applying a commercial whitener (Sunguard) to Granny Smith apples. The whitener had been developed for walnuts. They concluded from their experiments that Granny Smith apples could not be protected from sunburn by up to four topical applications of this particular whitening agent.

Miller Chemical & Fertilizer Corp. (Hanover, PA) markets an anti-transpirant concentrate called VAPOR GARD, and claims in its advertisements that the product reduced sunburn cullage by 30% in their trials. Transpiration is important to plant leaves, as evapotranspiration serves to cool the leaves and protects the leaves from heating to temperatures that are deleterious. Fruits have much lower transpiration rates than do leaves, but it seems likely that applying an anti-transpirant to fruit would exacerbate a situation in which there is already too much thermal energy.

Myhob, Guindy, and Salem in Egypt (*Bulletin of Faculty of Agriculture*, University of Cairo, 47(3):457-469 (1996)) reported that Agricultural GatCool significantly reduced sunburn as compared to controls sprayed with water on Balady mandarin fruits. duToit in South Africa (*Citrus and Subtropical Fruit Research Institute Information Bulletin No. 80*:8-9 (1979)) reported that spraying Koolcote on pineapple trees decreased fruit flesh temperatures by 2-3 degrees Celsius.

Lipton and Matoba (*HortScience* 6(4):343-345 (1971)) reduced sunburn of 'Crenshaw' melons by whitewashing fruit with a suspension of aluminum silicate.

Ing (*Good Fruit Grower* 49(6):58 (1998)), commenting on unpublished field trials, reports that the application of kaolin to apple fruits not only acts as an insect repellent, but also lowers canopy temperature, increases fruit size, and may reduce sunburn. However, as noted by Ing, application of kaolin to fruit surfaces is problematic. To achieve an insecticidal result, large amounts of kaolin (50 to 100

pounds per acre) must be applied to the fruit trees. Current kaolin formulations are reported to suffer from substantial application problems such as excessive foaming and "globbing" in spray tanks. (*Good Fruit Grower* 49(6):58 (1998)). Furthermore, kaolin powders are easily washed off by rain, thus necessitating multiple applications
5 in order to maintain beneficial effects. (*Good Fruit Grower* 49(6):58 (1998); *see also Washington State University Cooperative Extension Area Wide IPM Update* 3(4):1 (1998)).

Sekutowski et al. (U.S. Pat. No. 5,908,708) developed a protective water resistant coating that was formulated as an aqueous dispersion of particulate matter
10 having a hydrophobic outer surface in a low boiling point organic liquid, such as methanol. The particulate matter of the Sekutowski et al. coating can be any finely divided hydrophobic particulate solids including minerals, such as calcium carbonate, mica, talc, kaolin, bentonites, clays attapulgite, pyrophyllite, wollastonite, silica, feldspar, sand, quartz, chalk, limestone, precipitated calcium carbonate,
15 diatomaceous earth and barytes. One agricultural use of the Sekutowski et al. aqueous dispersions is to provide tree leaves with a water resistant coating by spraying the formulation onto the surface of the leaves. The water resistant coating is thought to reduce plant disease and insect damage. However, one major problem with the Sekutowski et al. formulation is the use of large volumes of organic liquids
20 such as alcohols, ketones and cyclic ethers that are highly flammable and pose other health risks to workers during spray application.

Protective formulations which additionally function as pesticides in plant crops would be a valuable addition to Integrated Pest Management (IPM) practices providing "soft" suppression of pests without disrupting natural control processes.
25 Desirable formulations would be expected to be non-toxic to mammals and thus safe for applicators and farm workers. Application of the protective formulations by commonly employed horticultural spray operations invariably involves treatment of foliage and fruit or vegetable. It is therefore important to develop new formulations that have protective properties against sunburn to fruits and vegetables as well as
30 against damage caused by insects that inhabit both foliage and fruit.

In summary, there is a lack of adequate means to prevent sunburn and insect damage to fruit and vegetable crops. Thus, there is a strong need in agricultural markets for an inexpensive and effective composition that prevents sunburn, repels deleterious insects, is long lasting, and is relatively amenable to easy application by
35 growers and commercial applicators.

Summary of the Invention

It has now been discovered that the foregoing problems can be overcome and that sunburn in apples, and other fruit and vegetable crops requiring exposure to high intensity solar irradiance for maturation, can be significantly reduced by treating the crop with an effective amount of a plant protective coating composition of the present invention. An effective amount of a plant protective coating composition of the invention is defined as any amount of the inventive composition that upon application to the surface of a fruit or vegetable, results in the measurable reduction of the incidence of fruit or vegetable sun damage. The plant protective coating compositions of the invention also forms a barrier that reduces insect inflicted damage to the fruit or vegetable.

In one aspect, the present invention provides a fruit or vegetable that is protectively coated with a plant protective composition comprising lipophilic thixotropic smectic clay suspended in a wax emulsion. The wax emulsion preferably comprises complex hydrocarbons (also known as a matrix of hydrocarbons), at least one emulsifying agent and water. In a presently preferred embodiment of the present invention, both an anionic lipophilic hydrophilic emulsifier and a cation hydrophilic emulsifier are used to emulsify the matrix of hydrocarbons. Preferably, the protective composition is a mixture of about 0.5 to 10% (weight/weight) of lipophilic thixotropic smectic clay dispersed in about 90 to 99.5% (weight/weight) of the wax emulsion. For some uses of the inventive composition it is preferable to dilute the mixed composition into an aqueous solution. Preferably, the compositions of the invention are diluted into an aqueous solution in a volume/volume ratio of between about 1 part plant protective composition to about 1 part aqueous solution to about 1 part plant protective composition to about 10 parts aqueous solution.

Preferred plant protective coating compositions are sprayable onto fruit trees, vegetable crops and the like by a wide variety of commercial agricultural applicators. The matrix of hydrocarbons helps to maintain the physical integrity of the clay film on the fruit surface making the formulation more durable and resistant to rain wash. Because the plant protective coating compositions, when applied as finely dispersed spray particles, cover both foliage and fruit, a dual beneficial effect is achieved through prevention of the incidence of sunburn and damage by insects. The physical integrity of the clay film, as well as the matrix of hydrocarbons on foliage and fruit surfaces also provide an effective protective barrier against harmful insects which may naturally reside on both foliage and fruit.

In the practice of the invention, proper dilution of the inventive composition in an aqueous solution allows effective spray application of the sun and insect protective material on to fruits or leaves prior to conditions that lead to the incidence of fruit sunburn or insect damage. The inventive composition is preferably sprayed
5 onto plants at a rate of 100 to 500 gallons per acre. As compared to other formulations and treatments used to prevent sunburn damage of fruits, the inventive compositions and methods of application significantly reduce the incidence of fruit sunburn damage resulting in both fruit necrosis and browning.

The inventive compositions and methods are applicable to a wide variety of
10 fruits and vegetables including, for example, apples, pears, tomatoes, peppers, curburbits, honeydew melons, cantaloupes, avocados, plums, beans, squashes, peaches, grapes, strawberries, raspberries, gooseberries, bananas, oranges, tulips, onions, cabbages, and other. See, for example, Brooks, C. and Fisher, D. F., "Some High-Temperature Effects in Apples: Contrasts in the Two Sides of an Apple," *J.*
15 *Agr. Res.* 32(1):1-16. (1926); Ware, W. M., "High Temperature Injury on the Growing Apple," *Gardners Chron.* 92:287-288 (1932); Meyer, A., "Comparative Temperatures of Apples," *Proc. Amer. Soc. Hort. Sci.* 28:566-567 (1932); Whittaker, E. C. and McDonald, S. L. D., "Prevention of Sunscald of Deciduous Fruit Trees in Hot Climates," *Agr. Gaz. N. S. Wales* 52:231-233 (1941); Moore, M. H. and Rogers,
20 W. S., "Sunscald of Fruits," *East Mallng Res. Sta. Report*, Pp. 50-53. (1943); Cook, M. T., "Sunburn and Tomato Fruit Rots," *Phytopathology* 11:379-380 (1921); Harvey, R. B., "Sunscald of Tomatoes," *Minn. Studies Plant Sci.* 4:229-234 (1924); Harvey, R. B., "Conditions for Heat Canker and Sunscald in Plants," *J. Forestry* 23:292-294 (1925); Ramsey, G. B. and Link, G. K. K., "Market Diseases of Fruits
25 and Vegetables: Tomatoes, Peppers and Eggplants," *U. S. Dept. Agr., Misc. Publ.* 121:28-29 (1932); Moore, M. H. and Rogers, W. S., "Sunscald of Fruits," *East Mallng Res. Sta. Report*, Pp. 50-53. (1943); Retig, N. and Kedar, N., "The Effect of Stage of Maturity on Heat Absorption and Sunscald of Detached Tomato Fruit," *Israel J. Agr. Res.* 17:77-83 (1967); Kedar, N. and Retig, N., "An Oblong Dwarf
30 Tomato Resists Sunscald," *New Scientist* 36:546 (1967); Weber, G. F., "Diseases of Peppers in Florida," *Florida Univ. Agr. Expt. Sta. Bull.* 244:35-37 (1932); Bremer, H., "On Pod Spots in Peppers," *Phytopathology* 35:283-287 (1945); Barber, H. N. and Sharpe, P. J. H., "Genetics and Physiology of Sunscald of Fruits," *Agr. Meterol* 8:178-191 (1971); Rabinowitch, H. D., Friedmann, M., and Ben-David, B.,
35 "Sunscald Damage in Attached and Detached Pepper and Cucumber Fruits at Various

- Stages of Maturity," *Scientia Hort.* 19:9-18 (1983); Rabinowitch, H. D., Ben-David, B., and Friedmann, M., "Light is Essential for Sunscald Induction in Cucumber and Pepper Fruits, Whereas Heat Conditioning Provides Protection," *Scientia Hort.* 29:21-29 (1986); Leclerg, E. L., "The Relation of Leaf Blight to Sun Scald of Honeydew Melons," *Phytopathology* 21:97-98 (1931); Lipton, W. J., "Ultraviolet Radiation as a Factor in Solar Injury and Vein Tract Browning of Cantaloupes," *J. Amer. Soc. Hort. Sci.* 102:32-36 (1977); Schroeder, C. A. and Kay, E., "Temperature Conditions and Tolerance of Avocado Fruit Tissue," *Calif. Avocado Soc. Yearbook* 45:87-92 (1961); Renquist, A. R., Hughes, H. G. and Rogoyski, M. K., "Solar Injury of Raspberry Fruit," *HortScience* 22:396-397 (1987); Maxie, E. C. and Claypool, L. L., "Heat Injury in Prunes," *Proc. Amer. Soc. Hort. Sci.* 69:116-121 (1956); Farmer, A., "Sunscald of Japanese Plum Fruits," *Orchardist New Zealand* 51:113-114 (1968); Macmillan, H. G., "Sunscald of Beans," *J. Agr. Res.* 13:647-650 (1918); Macmillan, H. G., "Cause of Sunscald of Beans," *Phytopathology* 13:376-380 (1923); Macmillan, H. G. and Byars, L. P., "Heat Injury to Beans in Colorado," *Phytopathology* 10:365-367 (1920); Ramsey, G. B. and Wiant, J. S., "Market Diseases of Fruits and Vegetables: Asparagus, Onions, Beans, Peas, Carrots, Celery, and Related Vegetables," *U. S. Dept. Agr., Misc. Publ.* 440:17-32. (1941); Ramsey, G. B., Wiant, J. S. and Link, G. K. K., "Market Diseases of Fruits and Vegetables: Crucifers and Cucurbits," *U. S. Dept. Agr., Misc. Publ.* 292:20 (1938); Rhoads, A. S., "Sun-scald of Grapes and its Relation to Summer Pruning," *Amer. Fruit Grower* 44:20-47 (1924); Graves, A. H., "Sunscald of Tulip Flowers," *Phytopathology* 27:731-734 (1937); Green, G. C., "The Banana Plant. In: The Effect of Weather and Climate Upon the Keeping Quality of Fruit," *World Meteorological Organization, Technical Note No. 53*:113-135 Geneva (1963); Wade, N. L., Kavanagy, E. E. and Tan, S. C., "Sunscald and Ultraviolet Light Injury of Banana Fruits," *J. Hort. Science* 68:409-419 (1993); Ketchie, D. O. and Ballard, A. L., "Environments Which Cause Heat Injury to Valencia Oranges," *Proc. Amer. Soc. Hort. Sci.* 93:166-172. (1968).
- In addition, the plant protective compositions can be used on trees whose foliage is susceptible to sunburn, such as maples, basswood, boxelder, black walnut, birch, balsam fir, Douglas fir, Eastern white pine and spruce as well as many fruit trees (Litzow, M. and Pellett, H., "Materials for Potential use in Sunscald Prevention," *J. Arboriculture* 9:35-38 (1983); Green, S. B., "Forestry in Minnesota," *Geological and Natural History Survey of Minnesota, St. Paul* 401 pp. (1902); Huberman, M. A., "Sunscald of Eastern White Pine, *Pinus Strobus* L.," *Ecology* 24:456-471 (1943)).

The inventive methods and compositions can also be used on plants that are not susceptible to sunburn but which are impacted by insect damage. In addition to the above listed plants that are susceptible to sunburn and insect damage, the following plants would independently benefit from the insect protective qualities of the
5 inventive plant protective composition: soybeans, potatoes, peas, lentils, apricots, cherries.

Detailed Description of the Preferred Embodiment

Two types of sunburn exist in apples. One is a lethal phenomenon that leads to a necrotic area on the fruit. Such fruit becomes cullage. This phenomenon occurs
10 when the sun-exposed side of apple skin reaches a temperature of $52^{\circ} \pm 1^{\circ}$ Celsius for only 10 minutes. The second type of sunburn is a sublethal phenomenon that leads to a browning of the apple skin (sometimes referred to as "buckskin"). These apples can be sold, but at a lower grade and price.

Solar light contains ultraviolet, visible, and infrared radiation. All fruits and
15 vegetables which develop a yellow or red coloration as part of their growth cycle require a certain quantity of ultraviolet and visible light to achieve the desired maturation color. Infrared light predominantly leads to excessive heating and associated damage to fruit surfaces. The plant protective compositions of the present invention selectively filter out the infrared portion of solar light but allow other light
20 components to pass. The inventive clay coating is therefore invisible to the unaided eye. In contrast, kaolin based formulations appear on the surface of sprayed fruits and leaves as a whitish-gray dust, which uniformly reflects all components of solar light, therefore depriving the developing fruit of the beneficial aspects of solar light.

In one aspect, the present invention provides a fruit or vegetable that is
25 protectively coated with a composition comprising lipophilic thixotropic smectic clay and a wax emulsion. The wax emulsion comprises a matrix of complex hydrocarbons, at least one emulsifier agent and water. Preferably, the wax emulsion contains two emulsifying agents: an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier. Preferably, each emulsifier is present in the wax emulsion at
30 a concentration of between 1-15% (weight/weight).

In another aspect, the present invention provides a method of protecting fruit and vegetables from sunburn, comprising treating a fruit or vegetable with a sunburn preventative amount of a plant protective composition comprising lipophilic thixotropic smectic clay and a wax emulsion. The wax emulsion is composed of a
35 matrix of complex hydrocarbons, at least one emulsifier agent and water. Preferably,

the composition is applied to the fruit or vegetable multiple times through the growing season.

5 In yet another embodiment of the invention a method of plant protection is provided, comprising treating a plant with an insect-controlling amount of a plant protective composition comprising lipophilic thixotropic smectic clay and a wax emulsion. The wax emulsion is composed of a matrix of complex hydrocarbons, at least one emulsifier agent and water.

10 The compositions and methods of the invention significantly decrease the incidence of both types of sunburn in apples. The plant protective compositions are preferably based on a thixotropic smectic clay material that is chemically altered to render its surface lipophilic. Thixotropic clays, in their original form are typically hydrophilic. In order to increase the ability of the protective compositions of the invention to adhere to the lipophilic surface of fruit, the clay is rendered lipophilic, such as, for example, by transformation by a chemical reaction of the clay with
15 quaternary ammonium compounds in which the ligands consist entirely of aliphatic long-chain hydrocarbons or of a mixture of aliphatic and aromatic hydrocarbon residues. This reaction converts the hydrophilic clay into a hydrophobic and lipophilic material that is capable of molecularly dispersing oils, waxes and other lipid-like materials including organic solvents. Suitable thixotropic clay materials
20 for use in the practice of the invention include clays that have been transformed by a chemical reaction of the clay with quaternary ammonium compounds and have a clay structure that weakens when subjected to shear forces and increases in strength upon standing. Many thixotropic smectic clays suitable for use in the practice of the present invention are commercially available through a variety of vendors.

25 As used herein, the term "smectic clay" material refers to a Bentonite, platelet-type clay. When transformed to render it lipophilic, this clay may also be referred to as "organoclay".

The successful functioning of the inventive sunburn protectant requires a matrix consisting of complex hydrocarbons which renders the formulation sprayable
30 by commercial agricultural applicators, maintains the physical integrity of the clay on fruit and allows passage of visible solar radiation needed for fruit color formation but reflects undesired solar infrared light. The wax emulsion is formed by emulsifying natural or synthetic waxes with at least one emulsifying agent. Preferably, both an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier are used to emulsify
35 the matrix of hydrocarbons. The wax emulsion in the protective compositions of the

present invention is intended to replace and enhance the properties of the natural wax layer which exists on the surface of all fruits and vegetables.

As used herein, the term "matrix of complex hydrocarbons" refers to a lipid based matrix that is capable of absorbing and dispersing the lipophilic organoclay. Suitable complex hydrocarbons for use in the present invention include, for example, natural and synthetic waxes that are suitable for human consumption, with melting temperatures that are higher than the melting temperatures of the target fruit or vegetable waxes. In a presently particularly preferred embodiment, the complex hydrocarbons of the present application is Carnauba Wax of a tropical origin. It contains a mixture of true waxes with long chain fatty acids and long chain esters. The fatty acid composition is complex but well represented by the term "Carnauba Wax" (*Corypha cerifera*). It will be apparent to those skilled in the art that other edible plant-derived waxes, such as Candelilla Wax (*Euphorbia cerifera* and *Pedilantus pavonis*), Alfa (*Stipa Tenacissima*), or mixtures thereof, will also be useful for this purpose. In addition, other natural wax mixtures well known in the art, such as montan wax, rice-bran wax, beeswax, Japan wax and mixtures thereof can also be used in the plant protective compositions of the present invention. It is also apparent that any edible synthetic waxes containing oxygen can also be used to practice the present invention. See, for example, the description of synthetic oxygen containing waxes in U.S. Patent No. 5,049,186, incorporated herein by reference.

The wax emulsion of the present invention is made by emulsifying the matrix of hydrocarbons with an amount of an emulsifying agent sufficient to emulsify the matrix of hydrocarbons. In this regard, a large number of different emulsifier agents can be used to prepare the wax emulsion used in the practice of the present invention. See for example the emulsifying agents described in U.S. Patent Nos. 5,049,186 and 5,165,915, incorporated herein by reference. Preferably, both an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier are mixed with the matrix of hydrocarbons in an amount sufficient to emulsify the edible waxes. Preferably, the anionic lipophilic and the ionic hydrophilic emulsifiers are each present in the wax emulsion at a concentration of between about 1-15% (weight/weight) relative to the matrix of hydrocarbons.

The anionic lipophilic surfactants employed in the practice of the invention have, preferably, a hydrophilic-lipophilic balance (HLB) ranging from 10 to 40. They are principally salts of fatty acids (for example alkaline salts or organic salts such as amine salts), the said fatty acids having, for example, from 12 to 18 carbon atoms,

and being able to have a double bond as in the case of oleic acid; the alkaline salts or salts of organic bases of alkyl-sulfuric and alkyl-sulfonic acids having 12 to 18 carbon atoms, of alkyl-arylsulfonic acids whose alkyl chain contains 6 to 16 carbon atoms, the aryl group being, for example, a phenyl group. They are also ether-sulfates, in particular, the sulfatation products of fatty alcohols and polyalkoxylated alkylphenols, in which the aliphatic chain has from 6 to 20 carbon atoms and the polyalkoxylated chain has from 1 to 30 oxyalkylene units, in particular oxyethylene, oxypropylene or oxybutylene. Preferred anionic hydrophilic surfactants are the fatty acids oleic acid and stearic acid.

Presently preferred ionic hydrophilic surfactants include amine compounds such as ethanolamine, diethanolamine, triethanolamine, alkyl alcohol amines such as methyl-ethanolamine, butyl-ethanolamine, morpholine and mixtures thereof.

The presently preferred wax emulsion for use as the wax emulsion in the plant protective coating composition of the present invention is APL-BRITE 310 C produced by Solutec Corporation (Yakima, WA). Other commercially available material suitable for use in the inventive protective coating composition are: Decco 231 produced by Elf-Atochem North America (Philadelphia, PA); Johnson's H.S and Johnson 31 produced by S.C. Johnson Wax (Racine, WI); and Shield Brite AP50C and Carnauba Gold produced by Pace International LLC (Seattle, WA).

A presently preferred material which meets the requirements specified for a chemically altered thixotropic smectic clay is Tixogel® MP 100 that can be commercially obtained from Süd-Chemie Rheologicals, a division of United Catalysts Inc. of Louisville, KY. Tixogel® MP 100 is presently employed as an additive to a wide range of products including cosmetics, but not to our knowledge for any treatments of fruits or vegetables and not in combination with a matrix of complex hydrocarbons. A person with skill in the art will appreciate that many other organoclay materials having the required clay properties exist. Representative examples of useful clay materials include: numerous Tixogel and Optigel products, also produced by Süd-Chemie Rheologicals; the Bentone line of organoclays, obtainable from Rheox, Inc. (Highstown, NJ); organoclays produced by Southern Clay Products (Gonzales, TX) and, the Vistrol and Organotrol lines of organoclays, sold by CIMBAR Performance Minerals (Cartersville, GA). The distinguishing property of the thixotropic organoclays used in the present invention is that they must be lipophilic.

For proper formulation of the inventive compositions it is essential to effect an activation of the organoclay (Tixogel® MP 100) with the wax emulsion (APL-BRITE 310 C) prior to dilution with water. A mixture of about 0.5 to 7% (weight/weight) Tixogel® MP 100 in APL-BRITE 310 C can be made at room temperature by mechanical stirring, but above about 7% (weight/weight) the mixture will quickly turn into a solid gel. Preferably, the plant protective composition is a mixture of about 5% (weight/weight) of Tixogel® MP 100 in about 95% (weight/weight) APL-BRITE 310 C. The resulting protective coating material contains thixotropic clay suspended in a sprayable wax emulsion. The ratio of thixotropic smectic clay to wax emulsion may change if products other than Tixogel® MP 100 or APL-BRITE 310C are employed as the organoclay and wax emulsion, respectively.

More generally, the plant protective composition of the present invention is a mixture of about 0.5 to 10% (weight/weight) lipophilic thixotropic smectic clay dispersed in about 90 to 99.5% (weight/weight) of the wax emulsion. Preferably, the plant protective composition is a mixture of about 3% to 7% (weight/weight) lipophilic thixotropic smectic clay dispersed in about 97 to 93% (weight/weight) of the wax emulsion. Most preferably, plant protective composition is a mixture of about 5% (weight/weight) lipophilic thixotropic smectic clay dispersed in about 95% (weight/weight) of the wax emulsion.

The wax emulsion comprises about 5% to 10% (weight/weight) natural wax or edible synthetic oxygen containing wax, about 2% to 30% (weight/weight) emulsifying agent and about 60 to 93% (weight/weight) water. Preferably, the emulsifying agent comprises about 1 to 15% (weight/weight) anionic lipophilic emulsifier, such oleic acid, and about 1 to 15% (weight/weight) ionic hydrophilic emulsifier, such as morpholene. When the anionic lipophilic emulsifier is oleic acid and the ionic hydrophilic emulsifier is morpholene, it is most preferable that morpholene be used at a molar ratio, relative to oleic acid, that is larger than about 1.0. Most preferably, the wax emulsion comprises about 5 to 10% (weight/weight) natural wax selected from the group consisting of Carnauba wax, Candelilla wax, Alfa wax, montan wax, rice-bran wax, beeswax, Japan wax and mixtures thereof, about 2 to 7% (weight/weight) oleic acid, about 2 to 7% (weight/weight) morpholene and about 76 to 91% (weight/weight) water.

The plant protective coating composition can be applied directly onto plants or it may be diluted in an aqueous solution in any ratio which accommodates the

desired field spray technique. Suitable ratios for use of the present invention include, for example, dilution of the protective coating mixture into an aqueous solution in a volume/volume ratio of from about 1 part protective coating mixture to about 1 part aqueous solution to about 1 part protective coating mixture to 10 parts aqueous solution. In most applications for apple and pear fruit, the rate of spray volume ranges from 100 to 400 gal/acre. The number of spray applications per growing season is also variable but ranges from one application up to ten applications depending upon weather conditions. A person skilled in the art will appreciate that the above mentioned rates would be expected to change to a minimal degree if the inventive composition were applied to other fruits and vegetables, except that there would be a greater variation in final mixture/water ratios due to the specific requirements of agricultural crops involved, i.e. row crops, perennial trees, etc.

Example 1

The beneficial effects of a representative protective composition of the invention in decreasing both types of sunburn in field trials on 'Jonagold' apples are shown in Table 1. The composition was 5% w/w of Tixogel® MP100 in APL-BRITE 310 C (hereafter PFT-X). PFT-X was applied at full strength onto apple fruits. A single application of the protectant was made to 'Jonagold' apples at Wenatchee, Washington on July 14, 1997. At the time of application no sunburn was observed on developing fruit. There was only one severe heat spell of sufficient intensity to cause the majority of sunburn during the 1997 season. It occurred during the first week of August. On August 19, apples treated with PFT-X had significantly less ($P < 0.05$) sunburn necrosis and sunburn browning than did untreated control fruits. On September 10, sunburn necrosis was significantly lower in treated apples. The incidence of the necrosis type of sunburn was decreased by 66% on fruits treated with PFT-X in these field trials. The incidence of the surface browning type of sunburn ("buckskin") was decreased by 79%. Total sunburn was decreased by 73% in apples treated in accordance with the invention.

Table 1. Incidence of Sunburn Necrosis and Sunburn Browning as Influenced by PFT-X Formulation

Fruit Variety	Observation Date	Incidence of Necrosis		Incidence of Browning	
		Control	Treated	Control	Treated
'Jonagold'	14 July 97	0 ¹	0	0	0
	29 July 97	6.7	5.0	6.7	0
	19 Aug. 97	26.3	9.1*	17.5	3.6*
	10 Sept. 97	25.9	8.8*	6.9	0

¹ Each mean represents observations on 60 attached fruit that had been fully exposed to solar radiation for a daily duration of 3 hours before to 3 hours after solar noon. Controls received no application of the test formulation. Treated apples received one application of formulation.

*Denotes statistical significance of differences between control and treatment for each date as determined by a Yates-corrected z-test at the 0.05 level with n= 60.

Example 2

The beneficial effects of a representative protective composition of the invention in decreasing sunburn in field trials on 5-year-old 'Jonagold' apples are shown in Table 2. The PFT-X composition was as listed in Table 1, but the formulation was diluted 1:1 with water before application to trees. Treatments were applied to single tree plots replicated ten times in a completely randomized design in the Clayton Orchard near Orondo, WA. All treatments were applied with a handgun sprayer at approximately 150 pounds per square inch (psi) to near the point of drip, simulating a dilute spray of approximately 200 gallons/acre. For PFT-X, this provided 40 pounds of organoclay per acre and for Surround®, this provided 50 pounds of kaolin per acre. Each formulation was applied three times during the 1999 fruit growing season on July 7, August 4, and September 1. The control trees were sprayed with water on the same dates. For comparison, Surround®, a kaolin-based formulation containing proprietary surfactants and spreaders (marketed on a limited scale in 1999 by Engelhard Chemical Co., Iselin, NJ) was applied in the same manner to another group of trees. Surround® was formulated as suggested by the manufacturer using M-03, a proprietary Spreader/Sticker. 450 ml of M-03 was added to 50 lbs of kaolin clay (Engelhard Chemical M-97-009) that had previously been added to 100 gallons of water in a recirculating sprayer tank.

The sunburn data are presented in Table 2. The incidence of sunburn in all treatments was evaluated on August 31 by evaluating all fruit on each tree in the experiment. The percent of sunburn incidence for each tree was calculated. Both sunburn necrosis and sunburn browning were evaluated, but the incidence of sunburn necrosis was so low (<7% of total sunburn) that the two types were combined and analyzed statistically. Data were transformed using the angular or inverse sine transformation method (Steel and Torrie, Principles and Procedures of Statistics, McGraw-Hill Book Co., Inc., New York) prior to an analysis of variance.

Table 2. Incidence of Sunburn as Influenced by PFT-X.			
Fruit Variety	Incidence of Sunburn (%)		
	Control	Treated with PFT-X	Treated with Surround®
'Jonagold'	15.77	6.01**	15.26
**Denotes statistical significance of differences between control and PFT-X at the 0.01 level.			
Total number of fruit evaluated were 723, 649, and 557 for the control, PFT-X treated, and Surround®-treated apples, respectively.			

The data in Table 2 indicate that apples treated in accordance with the invention showed significantly less sunburn than apples treated with water or Surround®.

Example 3

The beneficial effects of a representative protective composition of the invention in decreasing sunburn in field trials on 3-year-old 'Cameo' apples are shown in Table 3. Sunburn damage was evaluated September 1. Other experimental details were the same as those in Example 2 except that trees were smaller, and two trees were included in each replication. The trees were in the Fleming Orchard near Orondo, WA.

Table 3. Incidence of sunburn as influenced by PFT-X Application			
	Incidence of Sunburn (%)		
Fruit Variety	Control	Treated with PFT-X	Treated with Surround®
'Cameo'	13.40	6.59**	13.85
**Denotes statistical significance of differences between control and PFT-X at the 0.01 level. Total number of fruit evaluated were 291, 260, and 258 for the control, PFT-X treated, and Surround®-treated apples, respectively.			

The incidence of sunburn in 'Cameo' apples was reduced significantly when treated with the inventive PFT-X formulation as compared to apples treated with water or Surround® (Table 3).

5

Example 4

The beneficial effects of a representative protective composition of the invention in decreasing sunburn in field trials on 9-year-old 'Fuji' apples are shown in Table 4. Sunburn damage was evaluated October 19. Other experimental details were the same as those in Example 2 except that a fourth application of formulations was made September 29. All fruit on two large branches of each tree were evaluated, as trees were much larger than those used in Examples 2 and 3. The trees were in the Fugachee Orchards near Pateros, WA.

10

Table 4. Incidence of sunburn as influenced by PFT-X Application			
	Incidence of Sunburn (%)		
Fruit Variety	Control	Treated with PFT-X	Treated with Surround®
'Fuji'	14.85	2.44**	8.59
**Denotes statistical significance between PFT-X and both control and Surround® at the 0.01 level. Total number of fruit evaluated were 485, 779, and 489 for the control, PFT-X treated, and Surround®-treated apples, respectively.			

15

The incidence of sunburn in 'Fuji' apples was reduced significantly when treated with the inventive PFT-X formulation as compared to apples treated with water or Surround® (Table 4).

Example 5

To evaluate the entomological efficacy of the inventive formulation PFT-X, a trial was conducted with 12-year-old 'Gala' apple trees at the Washington State University Tree Fruit Research & Extension Center, Wenatchee, WA. Control of codling moth (*Cydia pomonella* L.)(CM) during their second generation was evaluated. PFT-X treatments were applied to single tree plots replicated five times in a randomized complete block. PFT-X was applied with a handgun sprayer at 300 psi to the point of drip, simulating a dilute spray of approximately 400 gallons/acre. Three different PFT-X and Surround® application protocols were tested:

- 1) trees were sprayed with PFT-X or Surround® three times during the CM oviposition period (July 19 [1,000 degree day total], July 27 and August 4, 1999);
- 2) trees were sprayed with PFT-X or Surround® three times during the CM hatch period (August 12 [1,250 degree day total], August 18 and 25); and
- 3) trees were sprayed with PFT-X or Surround® six times (all dates) covering the CM oviposition and hatch periods. For all PFT-X and Surround® application protocols a sample of fruits was harvested and an evaluation of CM insect damage to the fruit was made on September 1 by visually inspecting fifty apples per replicate and recording the number of stings and entries.

Table 5. Codling Moth damage to apple fruit as influenced by applications of PFT-X or Surround® during oviposition, hatch, or oviposition + hatch.

Treatment	Rate (Form./100 gal)	Timing ¹	#/50 fruit		% total injury
			Stings	Entries	
Surround®	25 lbs	Oviposition	0.8a ²	3.0bc	7.6b
Surround®	25 lbs	Hatch	0.8a	4.0b	9.6b
Surround®	25 lbs	Oviposition+hatch	0.8a	2.0bc	5.6b
PFT-X	20 lbs	Oviposition	0.8a	2.6bc	6.8b
PFT-X	20 lbs	Hatch	1.2a	2.2bc	5.2b
PFT-X	20 lbs	Oviposition+hatch	1.4a	0.2c	3.2b
Untreated	NONE		0.8a	12.2a	26.0a

¹Application dates for Oviposition timing were Jul 19, Jul 27 and Aug 4 and for the Hatch timing were Aug 12, 18, and 25. Applications for the Oviposition+hatch timing included all six dates.

²Means in the same column followed by the same letter not significantly different (P=0.05, Duncan's new multiple range test).

Both the PFT-X and Surround® treatments significantly reduced CM injury relative to the untreated control (Table 5). There was no difference in the number of CM stings (shallow unsuccessful entries) across treatments. Most of the effect of the treatments with both PFT-X and with Surround® was observed in the reduction of successful entries into fruit. There was no observed advantage of timing, but when applications were made to both the oviposition and hatch periods, the level of fruit injury was slightly lower than when treatments were applied to either the oviposition or hatch period. The formulations of the present invention show promise as tools to manage codling moth, probably as supplements to other "soft" tactics such as mating disruption. These data and the data presented in Tables 1-4 demonstrate that the inventive composition has dual benefits when applied to fruit trees. The inventive composition is effective at significantly reducing the incidence of fruit sunburn and reducing fruit damage caused by codling moth.

Example 6

Some formulations cause phytotoxicity and others affect physiological processes such as photosynthesis when applied to trees. It has been shown that any unusual change in the overall bioenergetic status of the plant can be detected by a change in chlorophyll fluorescence (*See generally*, Lichtenthaler, K.K., "Applications of Chlorophyll Fluorescence in Photosynthesis Research, Stress Physiology," *Hydrobiology and Remote Sensing*, Kluwer Academic Publishers, Dordrecht, Germany (1988)). This includes all the reactions from the oxidation of water through electron transport, development of the electrochemical gradient, ATP synthesis, and eventually the series of enzymatic reactions for CO₂ reduction to carbohydrate in the leaf. Even changes in the plant that affect stoma opening and gas exchange with the atmosphere are reflected by changes in the fluorescence characteristics of a leaf. Therefore fluorescence was used as an indicator of any deleterious effects resulting from application of formulation. An OS5-FL Modulated Chlorophyll Fluorometer (Opti-Sciences, Inc. Tyngsboro, MA) was used to determine 'dark-adapted' Fv/Fm. Fv/Fm = Fm - Fo/Fm where Fo and Fm are the minimal and maximal fluorescence yield of a 'dark adapted' sample. Fluorescence was determined on five attached leaves on trees in each of the five replications used in Example 4. On average, 84% of the incident quanta are absorbed by a leaf. Thus, a value for Fv/Fm of about 0.8 indicates healthy leaves with near maximal electron transport.

Table 6. Influence of PFT-X and Surround® on fluorescence of leaves (estimation of electron flow in Photosystem II of photosynthesis). Same trees and treatments used in Example 4 were tested.

Treatment	Rate of (Form./100 gal)	Application Dates	Fluorescence (Fv/Fm)
Surround®	25 lbs	Jul 19, Jul 27, Aug 4	0.777
Surround®	25 lbs	Aug 12, 18, and 25	0.797
Surround®	25 lbs	Jul 19, 27; Aug 4, 12, 18, 25	0.816
PFT-X	20 lbs	Jul 19, Jul 27, Aug 4	0.808
PFT-X	20 lbs	Aug 12, 18, and 25	0.781
PFT-X	20 lbs	July 19, 27; Aug 4, 12, 18, 25	0.785
Untreated	NONE		0.801

The results in Table 6 indicate that the inventive formulation had no significant effect on ($P = 0.05$) fluorescence of the leaves to which formulation was applied. Thus, no evidence of damage to the overall bioenergetic status of the trees is seen with any of the formulations. No phytotoxicity to either fruit or leaves was observed with any formulations.

Example 7

Before field testing, entomologists sometimes conduct bioassays to determine the inherent toxicity of new formulations, changes in behavior of insects exposed to new formulations, and appropriate concentrations to apply. Accordingly, the inventive PFT-X formulation was used in two bioassays.

Adulticide bean disk bioassay. Leaf disks (2 cm diameter) were cut from untreated leaves of bean (*Phaseolus vulgaris* 'Henderson Bush'). Disks were floated with the abaxial (lower) surface up in a 3/4 ounce plastic portion cup filled with cotton and distilled water. Twenty adult twospotted spider mites (TSM), (*Tetranychus urticae* Koch) were transferred to the lower surface with a fine paintbrush. The leaf disks containing mites were treated with five concentrations of PFT-X or a distilled water check.

All cups containing the five replicates of each treatment were treated at the same time in a Potter Spray Tower equipped with the intermediate nozzle, and set to 6.5 psi. Two ml of the pesticide solution were placed in the reservoir, and sprayed

onto the disks. The mites were held in a growth chamber at $22\pm 2^{\circ}\text{C}$. Mites were evaluated variously from 24 h after treatment for response as described immediately below.

Category	Description
Alive	Moving without stimulation, or capable of moving >1 body length after gentle stimulation with brush.
Dead	No movement whatsoever, even after stimulation; or desiccated.
Moribund	Capable of producing some movement, especially twitching of legs, but unable to move > 1 body length after stimulation.
Runoff	Found in cotton or water surrounding leaf surface, but not on leaf disk. Makes no difference if dead or alive. (If walk off occurs during the course of the evaluation, count as alive.)

5

Table 7 presents the results obtained using the bean disk bioassay and PFT-X at a variety of application doses.

Table 7. Mortality and runoff resulting from treatment of twospotted spider mites on bean disks treated with PFT-X. PFT-X was applied June 29, 1999, and the evaluation was done June 30. The full-strength PFT-X as described in Table 1 was diluted in distilled water to provide concentrations ranging from 100 to 700 grams of PFT-X per liter.			
Concentration (g/liter)	No. Subjects	% Mortality	% Runoff
700	111	7.3	1.0
500	103	3.8	3.5
300	99	0.0	4.6
200	101	2.9	1.9
100	102	4.9	0.0
0	103	4.5	4.6

10

The results in Table 7 indicate that there was no dose response to the inventive PFT-X formulation after 24 h, either in terms of mortality or runoff.

Motile Stage Mortality and Behavior, Whole Plant Bioassay: Five leaves on each of six infested bean plants from the 1998 composite TSM colony were tagged. Prior to treatment, all motile stages were counted with a 5x-magnification headband (OptiVisor). Counts from the top and bottom side of the leaf were recorded separately. The same leaves were counted 24 h after treatment. Various concentrations of PFT-X were applied with a hand-pump-pressurized sprayer. The suspensions were kept under constant agitation during application. Five replicates were used for each treatment. Table 8 shows the data obtained from the whole plant bioassays with the inventive PFT-X formulation applied at a variety of concentrations. Primary data were analyzed using the General Linear Models Procedure of SAS (SAS 1988 (*Statistical Analysis Institute*, 1988; *SAS/Stat User's Guide*, Release 6.03 Edition; SAS Institute, Inc., Cary, NC)) using both a classification model (AOV) and numeric (regression).

Table 8. Location and mortality status of mites before and after treatment with the inventive formulation in a whole bean plant bioassay. PFT-X was diluted as described in Table 7, and applied June 30, 1999. Pre-treatment observations were made before application on June 30, and post-treatment observations were made on July 1, 1999. Means in the same column followed by the same letter not significantly different.

Concn in g/liter	Live				Dead	
	Total live mites/leaf	Total surface mites/leaf	Bottom surface mites/leaf	Top surface % mites	Top surface mites/leaf	Bottom surface mites/leaf
<u>Pretreatment</u>						
700	35.6a	5.8a	29.8a	17.2	--	--
500	33.6a	4.8a	28.8a	15.9	--	--
300	35.8a	8.4a	27.4a	22.2	--	--
200	35.6a	8.0a	27.6a	23.6	--	--
100	38.2a	9.8a	28.4a	30.4	--	--
0	29.0a	12.6a	16.4a	42.9	--	--
<u>Post-treatment</u>						
700	7.2a	2.4a	4.8a	28.7	3.8	3.8
500	11.4a	3.8a	7.6a	36.4	2.2	4.0
300	6.8a	1.8a	5.0a	25.0	4.0	4.2
200	14.6a	4.2a	10.4a	27.7	2.8	2.4
100	12.2a	3.2a	9.0a	22.5	2.6	5.4
0	14.0a	6.6a	7.4a	42.6	4.8	3.6

Although there was a considerable decrease in mite population after treatment with PFT-X, this decrease was not related to concentration. No differences among the various concentrations of PFT-X occurred in any of the variables measured or calculated (Table 8). In addition to mortality, the behavior of the mites (i.e., occupation of the upper versus lower surface of the leaf) was observed. Normally, the TSM preferentially occupy the lower leaf surface, and most of the webbing is found there. Treatment with the PFT-X did not alter this pattern (Table 8). The relationship between concentration and percentage occupancy on the upper leaf surface was analyzed by regression analyses, but no significant relationship existed

after the treatment (data not shown). In summary, PFT-X does not appear to affect either mortality or one aspect of behavior (leaf surface preference) of these mites.

Example 8

The effects of the inventive formulation (PFT-X) on phytophagous mites and their natural enemies were examined in an apple orchard. Four-year-old 'Oregon Spur Delicious' apples were used. Treatments were applied with an air-blast sprayer calibrated to deliver 100 gallons per acre. PFT-X treatments were applied August 4, 1999. The plot originally had no mite populations, so the orchard was seeded with twospotted mites (*Tetranychus urticae* Koch) from a greenhouse colony and later with European red mites (*Panonychus ulmi* Koch) from another orchard. In addition, the plot was sprayed with Asana® 0.66EC (DuPont Co., Wilmington, DE)(1 pint/acre) plus Lorsban® 50W (Dow Chemical, Midland, MI)(3 lbs/acre) to reduce codling moth populations in the plots. Post-treatment mite counts were taken every week until early fall. A sample of 20 leaves per plot was taken and kept cool during transportation to the laboratory. Mites were removed from the leaves with a leaf-brushing machine, and collected on a revolving sticky glass plate. Mites on the plate were counted with the aid of a stereoscopic microscope. Motile and egg stages of the pest mites European red mite, twospotted spider mite, and McDaniel spider mite (*Tetranychus mcdanieli* McGregor) were counted, along with motile and egg stages of the predatory mites *Typhlodromus occidentalis* (Nesbitt) and *Zetzellia mali* (Ewing). Motile stages only of apple rust mite, *Aculus schlechtendali* (Nalepa), were also counted. The eggs of twospotted spider mite and McDaniel mite could not be distinguished from one another, and were recorded as a single category (*Tetranychus* eggs).

Table 9 presents the phytophagous and predatory mite population data and the effects of spray applications of various formulations including the inventive PFT-X composition.

Table 9. Phytophagous and predatory mite populations before and after treatment with miticides and formulations.

Treatment	Rate/acre	Total tetranychids/leaf		
		Aug 2	Aug 11	Aug 17
PFT-X	10 lbs.	6.99a ¹	6.92a	20.51a
PFT-X	20 lbs.	7.75a	9.95a	10.04a
Surround®	25 lbs.	6.74a	23.01a	19.24a
Surround®	50 lbs.	13.51a	8.91a	22.13a
Orchex 796 ²	1%	9.09a	21.25a	6.70a
Pyramite® 60W ³ + Orchex 796	4.4 oz.+0.25%	8.14a	5.83a	11.89a
Check	---	7.16a	13.93a	29.98a
Treatment	Rate/acre	Total predatory mites/leaf		
		Aug 2	Aug 11	Aug 17
PFT-X	10 lbs.	0.13a ¹	0.13a	1.30a
PFT-X	20 lbs.	0.00a	3.59a	0.00a
Surround®	25 lbs.	0.10a	3.43a	0.29a
Surround®	50 lbs.	0.00a	0.04a	0.38a
Orchex 796	1%	0.00a	0.79a	0.75a
Pyramite® 60W + Orchex 796	4.4 oz.+0.25%	0.03a	1.04a	0.09a
Check	---	0.18a	0.09a	0.33a

¹ Data were analyzed using analysis of variance on each count date (PROC GLM; SAS Institute, 1988). Means were separated with the Waller-Duncan k-ratio t-test.

² Purchased from Exxon Company, U.S.A., Houston, TX.

³ Purchased from BASF Agricultural Products, Research Triangle Park, NC.

The mite populations consisted primarily of twospotted mites (71% overall) with some European red mite, and occasionally, some McDaniel mite forming a proportion of the population. The predatory mite population was primarily *T. occidentalis* (82% overall), with the remainder of the population comprised of

Z. mali. Populations began to rise in late July, and were at an appropriate level (3 to 8 mites/leaf) by early August. No statistical differences occurred among any of the treatments (including the untreated check) at any time during the course of the experiment, despite treatment means that ranged from 7 to 30 mites/leaf (Table 9).

5 Predatory mite populations were high but variable throughout the test. On the first post-treatment count date (Aug 11), the low rate of Surround® and the high rate of PFT-X had exceptionally high *T. occidentalis* populations (Table 9). This is especially notable since Asana®, a chemical known for its toxicity to predatory mites, was being sprayed at intervals. The use of Asana® compromised the test for predator toxicity, but there was no evidence that any of the materials were acutely
10 toxic to *T. occidentalis* and *Z. mali*.

An additional mite control variable, known as cumulative mite days (CMD) was calculated for the formulations indicated in Table 9. CMD was calculated for each formulation using the equation:

$$15 \quad \text{CMD} = \sum 0.5(\text{pop}_1 + \text{pop}_2)(\text{date}_1 - \text{date}_2),$$

where pop_1 is the population (total tetranychids/leaf) on date_1 and pop_2 is the population (total tetranychids/leaf on date_2).

CMD represents a time-weighted measurement of the populations. The CMD for Pyramite® +Orchex (CMD = 402) was lowest. The CMD was 423 for PFT-X (10 lbs./A), and 477 for PFT-X (20 lbs./A). The CMD for the check was 567. The CMD was 508 for Surround® (50 lbs./A) and 519 for Surround® (25 lbs./A). For Orchex 796, the CMD was 513. The CMD data above indicate that PFT-X seemed to provide some suppression of the leaf mite populations across the growing season.

In summary, the inventive formulation of PFT-X tested in Table 9 had no apparent toxicity on the mites or their predators. As expected, PFT-X did not cause mortality in the mites. However, it is particularly important that the inventive formulation does not kill the beneficial predators or repel them from the leaf's surface, as this result indicates that PFT-X will be useful in Integrated Pest Management (IPM). In IPM practices, a formulation is useful only if the formulation provides what is called "soft suppression" of pests. That is, the IPM formulation does not cause a significant disruption to the natural control processes by, for example, negatively impacting populations of beneficial organisms.

Example 9

35 The effects of several formulations on leafhopper nymphs in an apple orchard (cv. 'Braeburn') near Quincy, WA were examined. Four replicates were used where

each replicate consisted of three trees in a single row. Leafhopper nymphs were sampled by counting the nymphs on 20 leaves/tree. Populations were sampled weekly until the majority of the population had transformed to the adult stage. A single-spray program and a three-spray program were compared. The single-spray treatment and the first application of the three-spray program were applied on August 3, 1999, using a multiple tank air-blast sprayer calibrated to deliver 100 gallons/acre. The second and third sprays of the three-spray program were applied on August 12 and August 20, 1999. Table 10 presents the data obtained from this study.

10

Table 10. Leafhopper nymph populations before and after treatment with pesticides and formulations.

Treatment	Rate/acre	No. appl.	Leafhopper nymphs/leaf					
			July 29	Aug 6	Aug 9	Aug 16	Aug 23	Aug 31
PFT-X	20 lbs	1	3.89a ¹	1.99bcd	0.91c	3.86abc	3.55ab	1.10ab
PFT-X	20 lbs	3	3.54a	2.81bc	2.85a	3.49abc	3.40ab	1.21ab
Surround®	50 lbs	1	3.44a	1.86bcd	1.09bc	2.38bc	2.63ab	1.36a
Surround®	50 lbs	3	3.49a	1.41cd	1.08c	1.88c	2.01bc	0.31b
Orchex 796	1%	1	3.44a	3.28b	3.36a	5.01ab	4.15a	1.65a
Pyramite® 60W + Orchex 796	4.4 oz +0.25%	1	3.53a	1.34cd	2.46ab	5.09ab	3.73ab	1.44a
Provado® L6F ² +Sylgard 309 ³	6 fl oz +4 fl oz.	1	3.70a	0.61d	0.20c	1.18c	0.60c	0.94ab
Check	---	---	3.70a	6.11a	3.79a	6.28a	4.24a	1.85a

¹ Data were analyzed using analysis of variance on each count date (PROC GLM; SAS Institute, 1988). Means were separated with the Waller-Duncan k-ratio t-test. Means within columns not followed by the same letters are significantly different.

² Purchased from Bayer Corporation, Pittsburgh, PA.

³ Purchased from Wilfarm, L.L.C., Gladstone, MO.

The inventive PFT-X formulation (single application on August 3) provided suppression of nymphs through August 9, but thereafter the population mean was not different from the check (Table 10). With the three-spray program, PFT-X significantly suppressed nymph populations only on August 6, although the population means for the nymphs were always lower than the check. Only the standard (Provado + Sylgard) provided much knockdown and residual control.

Orchex 796, an oil used by some in IPM programs as a soft pesticide, was included in this test. It was different than the check only on August 6. Its suppression of nymph populations was therefore much like that of the inventive PFT-X formulation. Thus, the data presented in Table 10 indicate that the PFT-X formulation of the present invention can be used as a component of an integrated pest management program.

Example 10

The beneficial effects of a representative protective composition of the invention in decreasing damage by deleterious insects to foliage and fruit is tested in field trials on (A) apples [cv. 'Delicious', 'Golden Delicious', 'Fuji', 'Cameo', 'Jonagold' and 'Gala'] with the following target insects: codling moth, leafrollers, leafhoppers, spider mites, aphids, leafminers, true bugs (*Pentatomidae* and *Miridae*), cutworms, fruit worms, apple maggot, cherry fruit fly and San Jose scale; and on (B) pears [cv. 'Bartlett' and 'd'Anjou'] with the following target insects: pear psylla, true bugs, cutworms, spider mites, mealybug, and codling moth. Initial tests are conducted with high-pressure handgun spray equipment using a spray volume equivalent to 100 to 400 gal/acre. The results obtained allow determination of an activity profile for the inventive formulation on the target insects. Increasing concentrations of Tixogel® MP100 from 1 to 5% in APL-BRITE 310 C are used with aqueous dilutions of 1/2 to 1/10 strength to arrive at appropriate concentrations. Treatments are replicated three to six times in a randomized complete block design with single trees or small blocks of trees. An appropriate control consists of trees that receive no spray treatments. For entomological evaluations of pests on foliage, populations of insects such as mites, aphids, leafhoppers, pear psylla, and leafminers are evaluated pre-treatment and at intervals in the post-treatment period to determine efficacy. For pear psylla and other pests such as the codling moth, scale, and leafrollers, the level of injury to fruit is evaluated at three times during the growing season in each treatment by checking at least 25 fruit per tree (replicate).

While the preferred embodiment of the invention has been illustrated and described, it will be appreciated that various changes can be made therein without departing from the spirit and scope of the invention.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A fruit or vegetable coated with a plant protective coating comprising lipophilic thixotropic smectic clay and a wax emulsion.
2. The fruit or vegetable of Claim 1 wherein the wax emulsion comprises a matrix of complex hydrocarbons, an emulsifying agent and water.
3. The fruit or vegetable of Claim 2 wherein the emulsifying agent comprises an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier.
4. The fruit or vegetable of Claim 1 wherein the wax emulsion comprises an edible synthetic oxygen containing wax, an emulsifying agent and water.
5. The fruit or vegetable of Claim 1 wherein the plant protective coating comprises about 0.5 to 10% (weight/weight) lipophilic thixotropic smectic clay and about 90 to 99.5% wax emulsion.
6. The fruit or vegetable of Claim 3 wherein the matrix of complex hydrocarbons comprises a wax mixture comprising long chain fatty acids and long chain esters.
7. The fruit or vegetable of Claim 6 wherein the wax mixture is a natural wax selected from the group consisting of Carnauba wax, Candelilla wax, Alfa wax, montan wax, rice-bran wax, beeswax, Japan wax and mixtures thereof.
8. The fruit or vegetable of Claim 3 wherein the anionic lipophilic emulsifier is selected from the group consisting of oleic acid, stearic acid and mixtures thereof.
9. The fruit or vegetable of Claim 3 wherein the ionic hydrophilic emulsifier is selected from the group consisting of ethanolamine, diethanolamine, triethanolamine, alkyl alcohol amine, morpholine and mixtures thereof.
10. The fruit or vegetable of Claim 5 wherein the protective coating mixture is diluted into an aqueous solution in a volume/volume ratio of from about 1

part protective coating mixture to about 1 part aqueous solution to about 1 part protective coating mixture to 10 parts aqueous solution.

11. A plant protective composition comprising:
about 0.5 to 10% (weight/weight) lipophilic thixotropic smectic clay, and
about 90 to 99.5% (weight/weight) wax emulsion, said emulsion comprising:
about 5 to 10% (weight/weight) natural wax selected from the group
consisting of Carnauba wax, Candelilla wax, Alfa wax, montan wax, rice-bran wax,
beeswax, Japan wax and mixtures thereof;

about 1 to 15% (weight/weight) oleic acid;
about 1 to 15% (weight/weight) morpholene; and
about 60 to 93% water.

12. A method of protecting a plant from sunburn, comprising treating said plant with a sunburn preventative amount of a plant protective composition comprising lipophilic thixotropic smectic clay and a wax emulsion.

13. The method of Claim 12 wherein the treated plant is selected from the group consisting of apple, pear, tomato, pepper, curburbit, honeydew melon, cantaloupe, avocado, plum, bean, squash, peach, grape, strawberry, raspberry, gooseberry, banana, orange, tulip, onion, cabbage, maple tree, basswood tree, boxelder tree, black walnut tree, birch tree, balsam fir, Douglas fir, Eastern white pine and spruce.

14. The method of Claim 12 wherein the wax emulsion comprises a matrix of complex hydrocarbons, an emulsifying agent and water.

15. The method of Claim 12 wherein the wax emulsion comprises an edible synthetic oxygen containing wax, an emulsifying agent and water.

16. The method of Claim 12 wherein the plant protective composition comprises about 0.5 to 10% (weight/weight) lipophilic thixotropic smectic clay and about 90 to 99.5% of the wax emulsion.

17. The method of Claim 14 wherein the emulsifying agent comprises an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier.

18. The method of Claim 14 wherein the matrix of complex hydrocarbons comprises a wax mixture comprising long chain fatty acids and long chain fatty alcohol esters.

19. The method of Claim 17 wherein the anionic lipophilic surfactant is selected from the group consisting of oleic acid, stearic acid and mixtures thereof.

20. The method of Claim 17 wherein the ionic hydrophilic emulsifier is selected from the group consisting of ethanolamine, diethanolamine, triethanolamine, alkyl alcohol amine, morpholine and mixtures thereof.

21. The method of Claim 17 wherein the plant protective composition is diluted into an aqueous solution prior to treating the plant.

22. The method of Claim 18 wherein the wax mixture is a natural wax selected from the group consisting of Carnauba wax, Candelilla wax, Alfa wax, montan wax, rice-bran wax, beeswax, Japan wax and mixtures thereof.

23. The method of Claim 16 wherein the plant protective composition is diluted into an aqueous solution in a volume/volume ratio of from about 1 part protective coating mixture to about 1 part aqueous solution to about 1 part protective coating mixture to 10 parts aqueous solution.

24. The method of Claim 12 wherein the plant is treated by spraying the composition onto the surface of the plant.

25. The method of Claim 24 wherein the composition is sprayed with an application rate of about 100 to 500 gallons per acre.

26. The method of Claim 24 wherein the composition is sprayed onto the plant multiple times.

27. A method of protecting a plant from insect damage comprising treating a plant with an insect-controlling amount of a plant protective composition comprising lipophilic thixotropic smectic clay and a wax emulsion.

28. The method of Claim 27 wherein the treated plant is selected from the group consisting of apple, pear, tomato, pepper, curburbit, honeydew melon,

cantaloupe, avocado, plum, bean, squash, peach, grape, strawberry, raspberry, gooseberry, banana, orange, tulip, onion, cabbage, potato, pea, lentil, apricot, cherry, onion, maple tree, basswood tree, boxelder tree, black walnut tree, birch tree, balsam fir, Douglas fir, Eastern white pine and spruce.

29. The method of Claim 27 wherein the wax emulsion comprises a matrix of complex hydrocarbons, an emulsifying agent and water.

30. The method of Claim 27 wherein the wax emulsion comprises an edible synthetic oxygen containing wax, an emulsifying agent and water.

31. The method of Claim 27 wherein the plant protective composition comprises about 0.5 to 10% (weight/weight) lipophilic thixotropic smectic clay and about 90 to 99.5% wax emulsion.

32. The method of Claim 29 wherein the emulsifying agent comprises an anionic lipophilic emulsifier and an ionic hydrophilic emulsifier.

33. The method of Claim 29 wherein the matrix of complex hydrocarbons comprises a wax mixture comprising long chain fatty acids and long chain fatty alcohol esters.

34. The method of Claim 32 wherein the anionic lipophilic surfactant is selected from the group consisting of oleic acid, stearic acid and mixtures thereof.

35. The method of Claim 32 wherein the ionic hydrophilic emulsifier is selected from the group consisting of ethanolamine, diethanolamine, triethanolamine, alkyl alcohol amine, morpholine and mixtures thereof.

36. The method of Claim 32 wherein the plant protective composition is diluted into an aqueous solution prior to treating the plant.

37. The method of Claim 33 wherein the wax mixture is a natural wax selected from the group consisting of Carnauba wax, Candelilla wax, Alfa wax, montan wax, rice-bran wax, beeswax, Japan wax and mixtures thereof.

38. The method of Claim 31 wherein the plant protective composition is diluted into an aqueous solution in a volume/volume ratio of from about 1 part

protective coating mixture to about 1 part aqueous solution to about 1 part protective coating mixture to 10 parts aqueous solution.

39. The method of Claim 27 wherein the plant is treated by spraying the composition onto the surface of the plant.

40. The method of Claim 39 wherein the composition is sprayed with an application rate of about 100 to 500 gallons per acre.

41. The method of Claim 39 wherein the composition is sprayed onto the plant multiple times.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/25350**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : A23B 4/10, A23L 1/059; A01G 13/00

US CL : 47/ Digest 6 and Digest 11; 426/99, 102, 615

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 47/ Digest 6 and Digest 11; 426/99, 102, 615

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

West search terms clay, coating, wax, emulsion, trees, apples, sunburn

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2,198,991 A (DUTTON) 30 April 1940, page 1, col. 1, lines 1-5, page 2, col. 1, lines 15-54/	1-41
Y	JP 50-21382 A (YAMIGISHI) 22 July 1975, abstract.	20
Y	US 4,946,694 A (GUNNERSON et al.) 07 August 1990, abstract and col. 2, lines 15-54.	1-11

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

12 JANUARY 2000

Date of mailing of the international search report

04 FEB 2000Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

HELEN PRATT

Telephone No. (703) 308-1978

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/25350

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☐

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/25350

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-11, drawn to a fruit or vegetable coated with a protective coating.

Group II, claim(s) 12-26, drawn to a method of protecting a plant from sunburn.

Group III, claim(s) 27-41, drawn to a method of protecting a plant from insect damage.

The inventions listed as Groups I-III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Groups one does not require the special technical feature of coating a plant to protect it from sunburn with a particular composition of Group II and Group III does not require the special technical feature of using a particular amount of the protective composition to control insect damage.